



United Behavioral Health

Behavioral Clinical Policy: Complementary And Alternative Medicine (CAM) Treatments For Behavioral And Substance Use Disorders

Document Number: BH727CAM_0622

Effective Date: June 21, 2022

Table of Contents

- [Introduction](#)
- [Instructions for Use](#)
- [Benefit Considerations](#)
- [Description of Service](#)
- [Coverage Rationale](#)
- [Clinical Evidence](#)
- [Applicable Codes](#)
- [References](#)
- [Revision History](#)
- [Appendix](#)

INTRODUCTION

Behavioral Clinical Policies are a set of objective and evidence-based behavioral health criteria used by medical necessity plans to standardize coverage determinations, promote evidence-based practices, and support members' recovery, resiliency, and wellbeing for behavioral health benefit plans that are managed by Optum®¹.

INSTRUCTIONS FOR USE

This guideline is used to make coverage determinations as well as to inform discussions about evidence-based practices and discharge planning for behavioral health benefit plans managed by Optum. When deciding coverage, the member's specific benefits must be referenced.

All reviewers must first identify member eligibility, the member-specific benefit plan coverage, and any federal or state regulatory requirements that supersede the member's benefits prior to using this guideline. In the event that the requested service or procedure is limited or excluded from the benefit, is defined differently or there is otherwise a conflict between this guideline and the member's specific benefit, the member's specific benefit supersedes this guideline. Other clinical criteria may apply. Optum reserves the right, in its sole discretion, to modify its clinical criteria as necessary using the process described in *Clinical Criteria*.

This guideline is provided for informational purposes. It does not constitute medical advice.

Optum may also use tools developed by third parties that are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Optum may develop clinical criteria or adopt externally-developed clinical criteria that supersede this guideline when required to do so by contract or regulation.

¹ Optum is a brand used by United Behavioral Health and its affiliates.

BENEFIT CONSIDERATIONS

Before using this policy, please check the member-specific benefit plan document and any federal or state mandates, if applicable. This Clinical Policy is provided for informational purposes. It does not constitute medical advice.

Pre-Service Notification

Admissions to an inpatient, residential treatment center, intensive outpatient, or a partial hospital/day treatment program require pre-service notification. Notification of a scheduled admission must occur at least five (5) business days before admission. Notification of an unscheduled admission (including emergency admissions) should occur as soon as is reasonably possible. Benefits may be reduced if Optum is not notified of an admission to these levels of care. Check the member's specific benefit plan document for the applicable penalty and provision of a grace period before applying a penalty for failure to notify Optum as required.

DESCRIPTION OF SERVICES

According to the National Center for Complementary and Integrative Health (NCCIH, 2021), treatments that are "complementary" or "alternative" represent approaches developed outside of mainstream Western, or conventional, medicine. These terms are often used interchangeably, but refer to different concepts:

- If a non-mainstream practice is used together with conventional medicine, it is considered "complementary";
- If a non-mainstream practice is used in place of conventional medicine, it is considered "alternative."

Acupuncture

Acupuncture describes varying procedures and techniques that involve the stimulation of points on the body. The most studied technique is insertion into the skin with thin, solid, metallic needles that are manipulated by either hands or electrical stimulation. A routine treatment may require 15 to 30 needles with follow-up treatments at 2-week intervals. Most commonly, acupuncture is used for back and neck pain, osteoarthritis, and headache. Research has also been conducted on the use of acupuncture to treat behavioral health conditions, such as depression and substance use disorder (Fowler, 2020).

Art Therapy

According to the American Art Therapy Association (AATA, 2017), art therapy is used to improve cognitive and sensorimotor functions, foster self-esteem, and self-awareness, cultivate emotional resilience, promote insight, enhance social skills, reduce, and resolve conflicts, and advance societal and ecological change. Art therapy combines the knowledge and understanding of human development and psychological theories/techniques with visual arts and the creative process. Art therapists incorporate the use of art media and verbal processing of produced imagery to help clients improve psychological health, cognitive abilities, and sensory-motor functions.

Dance/Movement Therapy (DMT)

DMT is defined as the psychotherapeutic use of movement to further the emotional, cognitive, physical, and social integration of the individual (American Dance Therapy Association [ADTA], 2020). Dance/movement therapy interventions apply affective, behavioral, motoric, cognitive, and systemic strategies, including the principles of development, wellness, and pathology. The use of specific methods, techniques, modalities, and verbal interventions within the practice of professional dance/movement therapy is restricted to professional dance/movement therapists appropriately trained in the use of such methods, techniques, or modalities. Dance/movement therapy may be identified by other terms in the research literature, including "dance movement psychotherapy", "dance therapy", "body psychotherapy", or "therapeutic movement".

Equine Therapy

Equine therapy uses the purposeful manipulation of equine movement to engage sensory, neuromotor, and cognitive systems in achieving functional outcomes (American Hippotherapy Association, 2020). Equine therapy can be conducted by physical therapists or occupational therapists

as part of a larger plan of care involving other neuro/sensorimotor techniques. Individual riding centers may also employ “certified path instructors” or “horsemanship instructors”. Equine therapy is identified by other terms in the research literature, including “hippotherapy”, “therapeutic horseback riding”, “horse therapy”, “therapeutic horsemanship”, and “equine-assisted therapy”. Behavioral health conditions for which riding centers promote their services include autism spectrum disorders, attention deficit hyperactivity disorder, post-traumatic stress disorder, and learning disability.

Music Therapy

Music therapy is the clinical use of music interventions to achieve individualized goals within a therapeutic relationship, and is typically conducted by an individual completing an approved music therapy program. Therapists may assess emotional well-being and social functioning through musical responses, and develop music sessions based on specific client needs. According to the American Music Therapy Association (AMTA), music therapy allows exploration of personal feelings and promotes positive changes in mood and emotional states (AMTA, 2022).

Naturopathic Detoxification

Naturopathic detoxification therapy (also known as “All-Natural Detox Therapy”, “Natural IV Therapy”, “Nicotinamide Adenine Dinucleotide (NAD) IV Therapy”, “Amino Acid Therapy”, “Neurotransmitter Restoration Therapy”, “Brain Restoration+”, “Gentle Detox”, “Easy Detox”, etc.) is part of a holistic approach to alcohol and drug addiction treatment. It involves an unknown and non-FDA-approved combination of vitamins, minerals, amino acids, and/or NAD coenzymes, administered intravenously and/or orally. This therapy claims to eliminate cravings from a drug or alcohol addiction and promote recovery. While the actual treatment regimen may vary by site, the following have been identified as common components of a naturopathic detoxification for substance abuse:

- Preadmission assessments, including a medical evaluation
- Laboratory testing to help determine individual need of the patient
- Approximately 10-15 IV infusions in addition to oral therapy – the content of the infusions and oral therapies is unknown (Miller et al., 2012).

Sauna/Niacin Detoxification

Sauna/niacin detoxification for substance use disorders (also known as “New Life Detoxification”, “sauna detoxification”, “Purification Rundown/Program”, “Purif”, “Effective Purification Program”, etc.) typically follows a protocol where the following components are delivered on a daily basis):

- Physical exercise
- Sauna, done in 30 minute sessions for up to 5 hours daily
- A multivitamin cocktail, the main ingredient of which is niacin
- Mineral supplements, including calcium, magnesium, iron, zinc, manganese, copper, iodine, and potassium

Treatment programs may be delivered at varying levels of care, depending on the individual patient. The purpose of sauna/niacin detoxification is to eliminate from the body any drug residues and other toxic substances that remain locked in fatty tissues and may be present in the blood stream (Lennox & Cecchini-Sternquist, 2018).

COVERAGE RATIONALE

The following complementary and alternative medicine treatments are unproven and not medically necessary for treating behavioral and substance use disorders due to insufficient evidence of efficacy:

- Acupuncture
- Art therapy
- Dance/movement therapy
- Equine therapy
- Music therapy
- Naturopathic detoxification
- Sauna/niacin detoxification (e.g., New Life Detox)

Summary of Clinical Evidence

Acupuncture

Armour et al. (2019) examined the effectiveness of acupuncture in major depressive disorder with a systematic review and meta-analysis of available literature. The goal of this study was to review the effectiveness of acupuncture compared to usual care (treatment as usual), compared to sham or placebo acupuncture, to a psychological intervention, and as an adjunct to selective serotonin reuptake inhibitors (SSRI) or serotonin and norepinephrine reuptake inhibitors (SNRI) medication. There were 29 eligible studies with 2268 participants; 22 trials were completed in China with 7 completed outside of China. Sample sizes ranged from 19 to 755. Participants needed to have depression as the primary diagnosis, be of any gender and of any ethnicity, aged 16 years or above, with clinically diagnosed depression. Outcome measures needed to include either a validated clinician rated measure of depression severity (such as the Hamilton Rating Scale for Depression (HAM-D)) or a participant-reported measure, such as the Beck Depression Inventory (BDI) or Patient Health Questionnaire (PHQ-9). The results showed that acupuncture produced clinically meaningful reductions in depression severity when compared to usual care (Hedges (g) = 0.41, 95% confidence interval (CI) 0.18 to 0.63), sham acupuncture (g = 0.55, 95% CI 0.31 to 0.79), and as an adjunct to antidepressant medication (g = 0.84, 95% CI 0.61 to 1.07). A significant relationship was found regarding an increase in the number of acupuncture treatments delivered and reduction in the severity of depression ($p = 0.015$). The authors report limitations such as the high risk of bias for participant blinding in the majority of studies; the relevance of the findings in Chinese populations compared to other populations is ambiguous, due to greater treatment frequency and total number of treatments in China. Lastly, the majority of trials did not report follow-up or durability data, and safety reporting was deficient. Future studies should strive to include durability follow-up data to determine if any acupuncture benefits and the severity of depression are sustained.

Li et al. (2019) conducted a systematic review of 10 meta-analyses and randomized controlled trials (RCTs) that met inclusion criteria regarding acupuncture as a treatment for anxiety or anxiety disorders. Participant sizes were small, ranging from 3-14. Four of the meta-analyses reported pooled results of acupuncture treatment (AT) versus sham acupuncture. The authors report that these pooled results show that the effect of acupuncture on anxiety is debatable. A meta-analysis of 6 RCTs compared acupuncture to placebo/sham acupuncture using the State-Trait Anxiety Inventory (STAI) score, they found a clinically irrelevant and non-significant reduction (mean differences (MD) = -1.54, 95% CI -4.73 to 1.64). All of the studies acknowledge that there is currently insufficient evidence due to the small size of the included trials or their low quality study design. Future high-quality randomized trials are needed to determine whether acupuncture efficacy is greater than other anxiety treatments.

Smith et al. (2018) conducted a Cochrane Review to examine the effectiveness and adverse effects of acupuncture in the treatment of depression. The review was part of an update to a previous Cochrane Review, and now contains data from 64 studies (7104 participants). Studies were included if they were randomized controlled trials comparing acupuncture versus control acupuncture, no treatment, medication, other structured psychotherapies (cognitive-behavioral therapy, psychotherapy, or counselling), or standard care. Treatment modalities included acupuncture, electro-acupuncture, and laser acupuncture. Most studies were at high risk of performance bias, at high or unclear risk of detection bias, and at low or unclear risk of selection bias, attrition bias, reporting bias, and other bias. The authors concluded that the reduction in severity of depression was less when acupuncture was compared with control acupuncture than when acupuncture was compared with no treatment control, although in both cases, results were rated as providing low-quality evidence. The decrease in severity of depression with acupuncture given alone or in conjunction with medication versus medication alone is uncertain owing to the very low quality of evidence. Few studies included follow-up periods or assessed important outcomes such as quality of life. According to the authors, high-quality randomized controlled trials are needed to examine the clinical efficacy and suitability of acupuncture, as well as its effectiveness, compared with acupuncture controls, medication, or psychological therapies.

van den Noort et al. (2018) conducted a systematic review to evaluate the use of acupuncture as an add-on treatment for patients with schizophrenia with a special focus on the treatment of accompanying sleep disorders. A total of 26 eligible studies with 1181 patients with schizophrenia who received acupuncture treatment were included in the review. The authors found that there is inadequate evidence for the use of acupuncture as add-on therapy in the treatment of patients with schizophrenia; however, positive results were found in the treatment of concomitant sleep disorders. This result needs to be confirmed in large, randomized, controlled trials.

Chen et al. (2018) conducted a systematic review and meta-analysis to assess the efficacy of acupuncture in treating opioid use disorder (OUD). A total of 9 studies with 1063 participants were included in the review. The results showed that acupuncture could be more beneficial than no treatment/sham acupuncture in terms of changes in craving for opioid, insomnia, and depression. There was insufficient evidence to support better outcomes with acupuncture compared to medication. In addition, these findings revealed that, compared to sham electroacupuncture (EA), EA had differences in alleviating symptoms of craving and depression. The authors concluded that acupuncture could be effective in treating OUD. Moreover, EA could effectively alleviate symptoms of craving for opioid and depression. Nevertheless, the conclusions were limited due to the low-quality and small number of included studies.

Amorim et al. (2018) reviewed the literature on the effectiveness of acupuncture and electroacupuncture for the treatment of patients with anxiety disorders in order to assess the scientific evidence for its use. The systematic review of the clinical research was focused on published clinical trials (controlled, randomized, and non-randomized) involving the treatment of anxiety with acupuncture. Only clinical trials where anxiety was treated as the therapeutic target, and not as a secondary measurement or being associated with other health condition or disease, were considered. Thirteen studies were identified to match exclusion and inclusion criteria and were selected for the analysis. Methodology, design, and quality of the research were highly variable. The authors concluded that there is good scientific evidence encouraging acupuncture therapy to treat anxiety disorders, however, additional research with robust study design and methodology is required confirm efficacy.

Hayes, Inc. (2021) completed a comparative analysis of 47 randomized controlled trials (RCTs) regarding acupuncture for the treatment of substance use disorder. The overall quality of evidence was identified as low. Current research shows a potential but unproven benefit. Certain published evidence suggests that safety and impact on health outcomes are at least comparable to standard treatment/testing. However, it remains unclear about safety and/or impact on health outcomes due to poor-quality studies, sparse data, conflicting study results, and applicability in general practice.

Gol et al. (2021) performed a double-blind, three-arm, randomized clinical trial to examine the additive effects of acupuncture to oral selective serotonin reuptake inhibitors (SSRIs) in decreasing anxiety symptoms. Participants (n=105) were divided into 3 groups; SSRIs alone (drug group, n=35), SSRIs with sham acupuncture (control group, n=35) and SSRI with acupuncture (acupuncture group, n=35), and treated for 4 weeks. Among the notable inclusion criteria was a confirmed anxiety disorder, adults aged 18-60 years, and no history of extensive SSRI use. The oral SSRIs prescribed were sertraline, citalopram, and escitalopram. Acupuncture was administered for 20 minute sessions, 3 times per week for 4 weeks. The participants and the psychiatrist were blinded and without knowledge of groupings. Participants were administered the State-Trait Anxiety Inventory (STAI) questionnaire at study start and at day 28. Serum cortisol levels were also monitored before the study and at day 28. STAI total score findings at day 28 indicated significant differences across all 3 groups (p <0.001). The results revealed that SSRIs with acupuncture (acupuncture group) showed the largest improvement in anxiety symptoms per the STAI scores, when compared to the drug group and the control group. Serum cortisol levels were not significantly modified across the 3 groups at baseline and day 28. Although the results show promise, the authors concluded that future research is needed to establish durability beyond 4 weeks with larger sample sizes to enhance generalizability.

Krause et al. (2020) conducted a randomized controlled three-arm study of National Acupuncture

Detoxification Association (NADA) acupuncture efficacy in alcohol addiction treatment. Adult participants (n=72) in the three-arm study were randomized into three groups; a twenty, 30-minute NADA group, sham acupuncture sessions group, and no intervention group. Symptoms such as craving, depression, anxiety and autonomic control of the heart were monitored. Measurement tools for depression, alcohol use, and anxiety were utilized that include Beck's Depression Inventory, Obsessive Compulsive Drinking Scale, and State-Trait Anxiety Inventory. Testing and monitoring was documented at baseline, immediately post intervention (sham intervention or control period) and at 4 weeks follow-up. One year after the study conclusion, abstinence was assessed. Results indicated that craving, anxiety, and depression were unchanged in the NADA acupuncture group or the sham group (p=n.s.). The one-year abstinence rates were unchanged across the 3 groups. A significant finding was improvement of cardiac autonomic function in the NADA acupuncture group; this improvement was maintained 4 weeks after intervention completion. The researchers note limitations such as small sample size with limited generalizability, blinding participants only, and replication in a larger sample size remains unclear. Future research is needed to determine efficacy and durability of acupuncture protocols for psychiatric conditions.

Art Therapy

Dunphy et al. (2019) conducted a systematic review of studies on creative arts interventions for older adults experiencing depression that examined outcomes of four creative arts modalities (art, dance movement, drama, and music); with particular attention paid to processes documented as influencing change in each modality; and mechanisms considered to result from these processes. An analysis of 75 articles (17 art, 13 dance, 4 drama, and 41 music) indicated mostly significant quantitative or positive qualitative findings, particularly for interventions led by creative arts therapists. Art therapy studies were found to be of medium quality. The primary concerns in the quantitative studies include a small sample sizes that were not randomized or blinded, a general lack of generalizability, and a lack of rigorous efforts to ensure validity in the results. Very few of the studies included follow up. Issues in qualitative studies in art therapy also relate to a lack of rigor to ensure credible data analysis and insufficient reporting in data collection. The authors recommend further research to assess the use of creative art modalities for depression.

Abbing et al. (2018) conducted a systematic review to evaluate the effects of art therapy (AT) on anxiety symptom severity in adults. Three randomized controlled trials with 162 participants in total met the inclusion criteria. Participants were diagnosed with PTSD, students with exam anxiety, or prisoners with anxiety. All studies had a high risk of bias and small sample sizes. The authors concluded that there is limited high quality evidence assessing the effectiveness of AT on anxiety; further high quality studies are required.

Deshmukh et al. (2018) assessed the effects of art therapy as an adjunctive treatment for dementia compared with standard care and other non-pharmacological interventions in a systematic review. Two randomized controlled trials (n = 60) met the inclusion criteria and were included in the review. In both studies there were no distinct changes reported between the intervention group and the control group in the important outcome measures. According to GRADE ratings, the authors reported the quality of evidence for these outcome measures to be 'very low'. The authors concluded that there is insufficient evidence regarding the efficacy of art therapy for people with dementia.

Dance/Movement Therapy

Salihu and colleagues (2021) completed a meta-analysis investigating dance interventions on depression symptoms, anxiety, and stress in adults with and without musculoskeletal disorders. Articles for the final selection were 28 randomized controlled trials with a total of 2249 participants. The majority of participants were females, aged 18-85, some with underlying health conditions such as Parkinson's disease (17.9%). Dance movement therapy was the most studied technique. Most of the control groups among the studies used normal daily activities as the usual care with some using music, physical or other activities. Frequency of the dance interventions were from once per week to twice per week, 60 minute sessions, for a duration of 12 weeks. Communities and care facilities were the most utilized locations for the interventions. Depression, anxiety, and stress levels were recorded using a variety of standardized instruments such as Depression-Anxiety-Stress Scale, the Beck Depression Inventory, and the Zung Self-Rating Scale. The quality of the studies were described as fair (n=12) and good (n=16), while the quality of evidence ranged from very low to low. The yielded

results were statistically significant for reducing depressive symptoms (SMD = -0.69, 95% CI -0.91 to -0.35, $p < 0.001$), for anxiety (SMD = -0.99, 95% CI = -1.92 to -0.05, $p < 0.05$), and for stress (SMD = -1.0, 95% CI = -1.83 to -0.17, $p < 0.05$). Statistical significance was found implementing at least 150 minutes per week of dance therapy to decrease depressive symptoms (SMD = -0.72, 95% CI -0.20 to -0.25, $p < 0.01$). Although the meta-analysis found statistical significance, the authors note that results should be considered with caution due to various limitations across the trials. A remarkable limitation is that none of the included trials were described as high quality. Additionally, there were design issues such as small sample sizes, lack of blinding, different types of dance with varied protocols, and heterogeneity within the characteristics of the participants. Further high quality studies are recommended and needed to establish effective dance therapy protocols and outcomes.

Dunphy et al. (2019) conducted a systematic review of studies on creative arts interventions for older adults experiencing depression that examined outcomes of four creative arts modalities (art, dance movement, drama, and music); with particular attention paid to processes documented as contributing to change in each modality; and mechanisms considered to result from these processes. An analysis of 75 articles (17 art, 13 dance, 4 drama, and 41 music) indicated mostly significant quantitative or positive qualitative findings, particularly for interventions led by creative arts therapists. Dance movement studies were largely randomized controlled trials (RCTs). The quality of dance studies varied, with ratings evenly distributed from the lowest to highest using the PEDro tool scores. Very few of the studies included follow up. Quality issues for this treatment approach relate to the lack of actual dance movement therapy (DMT) interventions. The authors recommend further research to examine the use of creative art modalities for depression.

In a randomized controlled trial, Mastrominico et al. (2018) examined the effects of dance movement therapy (DMT) on empathy for adults with autism spectrum disorder (ASD). The study was conducted as a multicenter study within the context of the EU-funded research project TESIS (Toward an Embodied Science of Intersubjectivity), and employed a two-factorial between-subject design. The treatment group ($n = 35$) participated in a 10-week manualized DMT intervention, whereas the control group ($n = 22$) received treatment only after a waiting period. Empathy, measured with the Cognitive and Emotional Empathy Questionnaire (CEEQ), was the main variable of interest, analyzed by a repeated measures analysis of variance. In order to also include incomplete data cases, the authors used the expectation-maximization algorithm for missing data estimation. Findings of the study suggest no significant changes in overall empathy between groups. The authors recommend that future studies focus more attention to the role of relationship and should test settings in which either the therapist or a co-therapist is doing the mirroring with the participant.

Meekums and Karkou (2017) performed a systematic review with the objective of examining the effects of dance movement therapy on behavioral, social, cognitive, and emotional symptoms of people with dementia when compared to no treatment or standard treatment. Electronic searches and personal communication were utilized to identify 102 studies. Nineteen full papers were reviewed with zero meeting the inclusion criteria of RCTs, including cross-over design and cluster RCTs. The authors concluded that there was no evidence for or against dance movement therapy as a viable intervention for dementia. The authors recommend future well-designed studies with a large number of participants are needed to assess dance movement therapy efficacy for dementia treatment.

Equine Therapy

White et al. (2020) completed a systematic review regarding the effect of equine-assisted therapies (EAT) for children with attention deficit/hyperactivity disorder (ADHD). Inclusion criteria were studies with primary quantitative study designs, children with a formal diagnosis of ADHD, and EAT interventions. Ten studies met the inclusion criteria, with ages ranging from 6-14 years, and 118 subjects. Overall positive developments were identified in behavioral, psychological, and physical outcome measures following the participation in an EAT. However, due to methodological limitations, caution is advised when interpreting these findings. The authors concluded that while EAT may offer some positive benefits for children with ADHD, further well-designed robust research is required to confirm efficacy.

Trzmiel et al. (2019) conducted a systematic review and meta-analysis to examine the effectiveness of Equine-Assisted Activities and Therapies (EAAT) in autism spectrum disorder (ASD) patients. A

total of 15 studies with 390 participants (aged: 3-16 years) were included. The interaction between psychosocial functioning and EAAT was investigated in most studies. Improvement was reported in the following areas: socialization, engagement, maladaptive behaviors, and shorter reaction time in problem-solving circumstances after EAAT. The authors indicated that the majority of the available reports demonstrated high effectiveness of EAAT, especially with regard to improved social functioning. The authors address that it is impossible to draw universal conclusions due to the considerable discrepancies in therapeutic protocols and measurement instruments of the above mentioned studies. According to the authors, the two main limitations of the review are the following: a relatively small sample size, which increases the risk of a calculation error, and differences in research methodology, which greatly hinders the comparison of the results. The authors suggest that further longitudinal research studies with standard protocols and large sample groups are needed to confirm efficacy.

Hawkins et al. (2019) performed a systematic review to assess animal-assisted therapy (AAT) in the treatment of schizophrenia. Eligible studies were randomized controlled trials that had compared animal-assisted therapy, or other animal-assisted intervention, to a control group using any participants with a clinical diagnosis of schizophrenia or related disorder, including schizophreniform or schizoaffective disorders and regardless of age, gender, setting, or severity and duration of illness. There were 6 full text articles that met inclusion criteria with the total number of participants as 390. The mean ages ranged from 34 to 79 years. Primary outcomes were mental state and behavior, clinical global response, and quality of life and wellbeing. Five out of seven studies included symptoms as an outcome measure, with one reporting improvements in negative symptoms and one study reporting improvements in positive and emotional symptoms. The remaining studies reported no significant effects of AAT. Three studies included quality of life as an outcome measure but did not find any noteworthy effects. Two studies revealed improvements in various measures of self-view. The authors denote that rigorous, large-scale randomized controlled trials with long-term follow-up are needed to determine the significance of AAT for schizophrenia. There is potential for the treatment of negative symptoms and negative self-view, however, currently results remain inconclusive.

Lai et al. (2019) conducted a systematic review regarding the efficacy of animal-assisted therapy (AAT) for people with dementia. Inclusion criteria was randomized controlled trials (RCTs), cluster-randomized trials, and randomized cross-over trials that compared AAT versus no AAT, AAT using live animals versus alternatives such as robots or toys, or AAT versus any other active intervention. There were 9 RCTs with 305 participants reviewed. Results from 2 studies with 83 participants revealed that people with dementia who had AAT were perhaps slightly less depressed at treatment conclusion than people who had standard care or other interventions not related to animals. Evidence from 3 studies with 164 participants showed that people who received AAT had no clear difference in their quality of life compared to those who did not. The authors report there was no evidence of an effect on social functioning (interactions with their environment and families), behavior, agitation, activities of daily living, self-care ability or balance. There were no clear variances when AAT was compared with the use of a robotic animal in 2 studies with 156 participants (in social functioning, behavior, and quality of life), or with the use of a soft toy cat in 1 study with 64 participants (in social functioning). The authors report that AAT may potentially decrease depressive symptoms, however, there is no clear evidence on whether AAT is beneficial or safe for people with dementia. Further studies with robust design are required to determine benefits of AAT for dementia.

Srinivasan et al. (2018) conducted a focused systematic review to evaluate the effects of equine therapy in individuals with autism spectrum disorder. Inclusion criteria for studies peer-reviewed articles and/or studies reporting data on treatment effects of "equine" therapy using experimental or quasi-experimental study designs. This review included 15 studies with 428 participants. The review suggested that equine therapy has beneficial effects on behavioral skills and to some extent on social communication in ASD. According to the authors, the evidence for positive effects of equine therapy on perceptuo-motor, cognitive, and functional skills is currently limited. Future studies are required, using rigorous study designs with large sample sizes to analyze the role of equine therapies as a treatment option for individuals with ASD.

Music Therapy

Icel and Basogul (2021) completed an experimental study to investigate the effects of progressive muscle relaxation training and music therapy on anger and sleep quality. Participants (n=66) were chronic psychiatric patients, ages 18-59, attending the Community Mental Health Center (CMHC). Participants were divided into a control group (n=32) and intervention group (n=34). Progressive muscle relaxation training with music therapy were implemented in the intervention group for 2 sessions per week, 1 hour sessions, for 3 months. The Pittsburgh Sleep Quality Index (PSQI) and the Trait Anger/Anger Expression Inventory were used for the pre and post-test measures. The overall findings showed statistical significance difference between the pre-test and post-test mean scores of the intervention group ($p \leq .001$). After the intervention, a statistically significant difference was indicated between the PSQI, anger-in, anger-out, anger-control, and trait anger scale scores of the two groups ($p \leq .001$). An additional notable result was a statistically significant relationship between sleep quality and anger scores in both the pre-test and post-test ($P < .05$); the post-test result reveals that the intervention controlled anger expression while sleep quality increased. According to the researchers, the results of this study support relaxation training and music therapy combined with pharmacology as beneficial to chronic psychiatric patients. Limitations of the study include small sample size, limited to CMHC participants with lack of generalizability, and lack of durability follow-up data. Future research with robust study design is recommended to further establish and support efficacy.

Rabeyron and colleagues (2020) compared music therapy and music listening for children and autism spectrum disorder in a randomized controlled trial. Thirty-seven participants aged 4 to 7 years old were recruited for this study from psychiatric facilities in France. The 37 participants were randomly assigned to either the Music Therapy (MT) group or the Music Listening (ML) group using a generated randomization list for each group at t0 (start point of study). The MT consisted of 25 sessions that lasted 30 minutes with a qualified music therapist; 19 participants completed all sessions. The ML consisted of 25 sessions that lasted 30 minutes with no specific therapeutic intervention; 17 participants completed all ML sessions. All sessions occurred between October 2014 and June 2015 at the same room in each facility on the same schedule every week, except during holidays. Outcomes were measured by The Clinical Global Impression (CGI) a 7-point scale, The Childhood Autism Rating Scale (CARS), a 15 -item scale, and The Aberrant Behavior Checklist (ABC), a 57-item checklist. The results showed CGI scores improved to a better extent in the MT ($d=2.16$) than in the ML condition ($d=1.33$) with a large effect size at t1 (the endpoint of the study), ($d=0.80$). This improvement was clinically significant as 63.2% of the children scored a decrease of at least 2 points on the CGI in the music therapy group compared to 29.4% in the music listening group. The ABC subscales noted significant improvement regarding lethargy and stereotypy symptoms. There were no clinically meaningful results for t0 and t1 and the CARS. The authors report a variety of limitations such as not relying on gold standard evaluation tools for autism spectrum disorder and no follow-up to test the durability of the music interventions. Future studies are encouraged to address these limitations and to include larger sample sizes.

In an updated Cochrane review, Geretsegger et al. (2017) evaluated the effects of music therapy, or music therapy added to standard care, compared with placebo therapy, standard care, or no treatment for people with serious mental disorders such as schizophrenia. Ten new studies were added to this update; 18 studies with a total 1215 participants are now included in the review. These studies examined effects of music therapy over the short, medium, and long-term, with treatment dosage varying from 7 to 240 sessions. Generally, most information is from studies at low or unclear risk of bias. The authors concluded that moderate to low-quality evidence suggests that music therapy as an addition to standard care improves the global state, mental state (including negative and general symptoms), social functioning, and quality of life of people with schizophrenia or schizophrenia-like diagnoses. However, effects were inconsistent across studies and depended on the number of music therapy sessions as well as the quality of the music therapy provided. The authors state that further research should focus on the long-term effects of music therapy, dose-response relationships, in addition to the relevance of outcome measures in relation to music therapy.

van der Steen et al. (2018) assessed the effects of music-based therapeutic interventions for people with dementia on emotional well-being including quality of life, mood disturbance or negative affect, behavioral problems, social behavior and cognition at the end of therapy and 4 or greater weeks after

the treatment conclusion. Twenty-two studies with 1097 randomized participants were included. Twenty-one studies with 890 participants contributed data to meta-analyses. Participants in the studies had dementia of varying degrees of severity, and all were resident in institutions. Seven studies delivered an individual music intervention; the other studies delivered the intervention to groups of participants. The methodological quality of the studies varied. All were at high risk of performance bias and some were at high risk of detection or other bias. The authors concluded that providing people with dementia who are in institutional care with a minimum of 5 sessions of a music-based therapeutic intervention probably reduces depressive symptoms and improves overall behavioral problems at the end of treatment. It potentially improves emotional well-being, quality of life, and decreased anxiety, however, may have little or no effect on agitation, aggression, or cognition. There is uncertainty about effects on social behavior and durability of this approach. According to the authors, future studies are needed to examine the duration of effects in relation to the overall duration of treatment and the number of sessions.

Geipel et al. (2018) systematically reviewed and quantified the effects of music-based interventions in reducing internalizing symptoms (i.e., depression and anxiety) in children and adolescents using a meta-analytical approach. Five studies were included and focused on the treatment of children and adolescents with either pathological depressive symptoms or pathological anxious symptoms. The participant ages ranged from 8-18 years. Analysis of data from randomized controlled trials, showed a significant main effect, indicating a greater reduction of internalizing symptoms in youth receiving music-based interventions (n = 100) compared to different control group interventions (n = 95). The existing evidence is limited to studies of low power and methodological quality. Included studies were highly heterogeneous with regards to the nature of the intervention, the measurements applied, the samples studied, and the study design. The authors concluded that study findings identify that music-based interventions may be efficient in reducing the severity of internalizing symptoms in children and adolescents. While these results are promising with the application of music-based intervention, more research adopting well controlled study designs of high methodological quality is needed.

A Cochrane review by Aalbers et al. (2017) analyzed the effects of music therapy for patients who were experiencing depression. Nine studies involving 421 subjects met the authors' inclusion criteria. Music therapy was found to provide positive short-term effects for depressive symptoms. However, no eligible evidence on functioning, adverse events or anxiety was identified. Future trials with larger participant sample sizes and robust study designs are necessary to confirm these findings.

Zhang and colleagues (2017) analyzed the efficacy of music therapy in elderly people diagnosed with dementia and whether music therapy can be used as first-line non-pharmacological treatment. A total of 34 studies were included with the publication dates from 1999 to 2015. The inclusion criteria were clinical trials with elderly people with dementia that compared any form and intervention method of music therapy with no music care. The authors excluded studies that did not provide comparative or missing outcomes. There were 16 studies identified as RCTs, 10 were controlled clinical trials (CCTs), and the other 8 were RCT/crossover trials. The primary outcomes were disruptive behavior and cognitive function; the secondary outcomes were depressive score, anxiety score, and quality of life. This meta-analysis indicated that music therapy had positive results for disruptive behavior and anxiety, with a positive trend for cognitive function, depression, and quality of life. Most trials revealed that music therapy was associated with an improvement in disruptive behavior and cognitive function outcomes. However, a majority of these associations did not reach statistical significance, and heterogeneity existed in most of the outcomes. The authors conclude that further clinical trials of music therapy should focus on large size samples, well-designed and randomized to confirm the effects of music therapy on disease specific outcomes, particularly in elderly people with dementia.

Naturopathic Detoxification

Miller and colleagues (2012) evaluated a natural dopaminergic agonist to improve dopaminergic function in substance use disorders. Participants were administered either oral-only treatment or IV treatment with Neuroadaptagen Amino Acid Therapy (NAAT) variant [KB220] along with other (oral) vitamin and mineral nutrients. The participants were polydrug abusers and in all cases drank alcohol to excess. The participants were detoxified from drugs within the last two months and had symptoms of craving behavior associated with protracted abstinence. The basic patented formula for NAAT Variant [KB220] included amino acid precursors such as L-phenylalanine, L-tyrosine, L-tryptophan, 5-

hydroxytryptophan, L-glutamine, a serotonin concentrating substance chromium, an enkephalinase inhibitor D- phenylalanine, a neurotransmitter synthesis promoter vitamin B6, as well as both methionine and leucine. The amounts of these ingredients varied according to individualized assessment. The IV administration was a 4-hour infusion once a day, over seven days. For the oral therapy protocol, everyone received nutrients including thiamine, riboflavin, niacin, B6, folate, B12, pantothenic acid, magnesium, choline, para-aminobenzoic acid, lecithin, and inositol. In addition, those who met the criteria for being serotonin deficient also received vitamins A, C, E, K, and D, glycine, leucine, DLPA, tyrosine, boron, calcium, biotin, zinc, potassium, methionine, selenium, copper, iodine, and manganese. Those who met the criteria for being dopamine deficient also received iodine, zinc, copper, selenium, manganese, chromium, potassium, boron, calcium, biotin, and 5-HTP. In the first phase of the study (n = 49) The authors determined that the IV and oral group did significantly better than the oral-only group over the first week and 30- day follow-up period on chronic symptoms, as measured by the Chronic Abstinence Symptom Severity (CASS) Scale. In the second phase of the study (n = 129), the combination of IV and oral treatment was provided to all subjects, and three factors (emotion, somatic, and impaired cognition) were extracted for baseline CASS-Revised variables. All three scales showed significant declines from pre- to post-treatment. In the third phase of the study, a total of 23 subjects were followed-up at six months, one year, and two years post-IV treatment via phone interview to determine both sobriety and relapse rates. A total of 21 (91%) reported being sober at six months with 19 (82%) having no relapse; 19 (82%) reported being sober at one year with 18 (78%) having no relapse; and 21 (91%) reporting being sober at two years post-treatment with 16 (70%) having no relapse. It is noted that the major limitation of the experiment was the small sample size. The authors recommend further research to confirm these results in a larger population and with the use of an accurate method of randomization.

Sauna/Niacin Detoxification

Hussain et al. (2018) examined the role of saunas, or whole-body thermotherapy as potential treatment for various health issues. Information was obtained using a voluntary online 71-item questionnaire on the individual characteristics, sauna-related habits, and perceived health and wellness experiences of regular sauna bathers. The study was conducted from October 2016 to October 2017. The validated 'SF-12' quality of life scoring tool was integrated into the questionnaire to measure physical and mental indicators of well-being. There were 482 valid responses recorded from around the world, with the age range of 17-80 years. Respondents sauna bathed approximately of 4-12 times each month (median of 6 times, n=443), which extrapolates to a frequency of approximately 1-2 occasions per week. Respondents reported one or more medically-diagnosed health conditions (32.1%, n=135/420). This study identified that sauna use has perceived health benefits that vary from relaxation, stress, relief, invigoration, and socializing to more specific health advantages such as aiding circulation, improving sleep, improving mental health, enhancing 'detoxification', and relieving back/musculoskeletal pain. The few reported incidences of adverse reactions to sauna bathing were mild. This study demonstrates that the use of a sauna for health purposes is not well established in the scientific community. The authors acknowledge numerous limitations of this study and recommend further research surrounding the use of saunas for improving health.

Lennox and Cecchini-Sternquist (2018) completed a prospective chart review of 109 individuals sequentially enrolled into the Hubbard sauna regimen as part of a multi-modal, long-term residential substance abuse treatment facility. The Hubbard regimen is based on exercise, sauna, and therapeutic nutrients. Data from medical charts, client self-reports and Short Form Health Survey (SF-36) responses indicated that the Hubbard sauna detoxification method was well tolerated, with a 99% completion rate, including 1 human immunodeficiency virus and 9 hepatitis C positive subjects. Statistically significant improvements were identified in both mental and physical SF-36 scores at regimen completion, in addition to the Addiction Severity Index and Global Appraisal of Individual Needs Short Screener change scores at rehabilitation program discharge, compared with enrollment. There were no serious medical complications, a very low discontinuation rate, and high participant satisfaction. The SF-36 results indicated improved physical and emotional symptoms. The authors recommend further research into this sauna-based treatment regimen. Future research should focus on additional outcomes measurements of physical and mental health changes with analysis of whether these are improved via toxic elimination, nutrient and systems restoration, or a combination of these methods.

Guidelines & Consensus Statements

Department of Veterans Affairs and Department of Defense (VA/DoD)

The Department of Veterans Affairs and Department of Defense (VA/DoD) Clinical Practice Guidelines for the Management of Major Depressive Disorder (2022) indicates the following for complementary and alternative treatments:

- For patients with major depressive disorder (MDD), there is insufficient evidence to recommend for or against acupuncture as an adjunctive treatment to pharmacotherapy
- For patients with MDD, there is insufficient evidence to recommend for or against yoga, tai chi, or qi gong as an adjunctive treatment to pharmacotherapy.

American College of Physicians (ACP)

In clinical guidelines on the nonpharmacologic versus pharmacologic treatment of adult patients with major depressive disorder, the ACP evaluated the use of complementary and alternative medicines (including acupuncture) and did not recommend their use (Qaseem et al., 2016).

U.S. FOOD AND DRUG ADMINISTRATION

As the practice of CAM has increased in the United States, the Food and Drug Administration (FDA) has seen increased confusion as to whether certain products used in CAM are subject to regulation under the Federal Food, Drug, and Cosmetic Act. See the following FDA website for more information: <http://www.fda.gov/RegulatoryInformation/Guidances/ucm144657.htm>.

Some niacin products are FDA-approved prescription products for treating high cholesterol; these prescription niacin products are typically available in high strengths. Both niacin and niacinamide are approved by the FDA for treatment and prevention of niacin deficiency, and certain conditions related to niacin deficiency. Niacin and other naturopathic therapy combinations have not been reviewed by the FDA for substance use detoxification.

CENTERS FOR MEDICARE AND MEDICAID SERVICES

Medicare does not have a National Coverage Determinations (NCDs) for the following complementary and alternative medicine modalities used in treating behavioral disorders and/or substance use:

- Art therapy
- Dance/movement therapy (DMT)
- Equine therapy
- Music therapy
- Naturopathic detoxification
- Sauna/niacin detoxification (also known as "New Life Detoxification", "sauna detoxification", "Purification Rundown/Program", "Purif", "Effective Purification Program", etc.)

Medicare does not cover acupuncture as an anesthetic or as an analgesic or for other therapeutic purposes. Refer to the following NCDs (www.CMS.gov):

- *NCD for Acupuncture (30.3)*
- *NCD for Acupuncture for Fibromyalgia (30.3.1)*
- *NCD for Acupuncture for Osteoarthritis (30.3.2)*

APPLICABLE CODES

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member-specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other clinical criteria may apply.

Procedure Codes	Description
97810	Acupuncture, 1 or more needles; without electrical stimulation, initial 15 minutes of personal one-on-one contact with the patient

97811	Acupuncture, 1 or more needles; without electrical stimulation, each additional 15 minutes of personal one-on-one contact with the patient, with re-insertion of needles(s). (List separately in addition to code for primary procedure.)
97813	Acupuncture, 1 or more needles; with electrical stimulation, initial 15 minutes of personal one-on-one contact with the patient
97814	Acupuncture, 1 or more needles; with electrical stimulation, each additional 15 minutes of personal one-on-one contact with the patient, with re-insertion of needles(s). (List separately in addition to code for primary procedure.)
90899	Unlisted psychiatric service or procedure
G0176	Activity therapy, such as music, dance, art or play therapies not for recreation, related to the care and treatment of patient's disabling mental health problems, per session (45 minutes or more)
H2032	Activity therapy, per 15 minutes
S8940	Equestrian/hippotherapy, per session

CPT® is a registered trademark of the American Medical Association

REFERENCES

1. Aalbers, S., Fusar-Poli, L., Freeman, R.E., Spreen, M., Ket, J.C.F., Vink, A.C., Maratos, A., Crawford, M., Chen, X.J., & Gold, C. (2017). Music therapy for depression. *Cochrane Database Systematic Review*, 11, CD004517, DOI: 10.1002/14651858.CD004517.pub3.
2. Abbing, A., Ponstein, A., van Hooren, S., de Sonnevile, L., Swaab, H., & Baars, E. (2018). The effectiveness of art therapy for anxiety in adults: A systematic review of randomised and non-randomised controlled trials. *PLoS One*, 13(12), e0208716.
3. American Art Therapy Association (AATA). (2017). What is art therapy? AATA website: <http://arttherapy.org/aata-aboutus/>.
4. American Dance Therapy Association (ADTA). (2020). What is dance/movement therapy? ADTA website: <https://www.adta.org/faq>.
5. American Hippotherapy Association, Inc (AHA). (2020). Frequently asked questions. AHA website: <https://www.americanhippotherapyassociation.org/what-is-hippotherapy>.
6. American Music Therapy Association (2022). Frequently asked questions. AMTA website: <http://www.musictherapy.org/faq/>.
7. American Psychiatric Association. (2022). *Diagnostic and statistical manual of mental disorders, (5th ed.), Text Revision*. American Psychiatric Publishing.
8. Amorim, D., Amado, J., Brito, I., Fiuza, S.M., Amorim, N., Costeira, C., & Machado, J. (2018). Acupuncture and electroacupuncture for anxiety disorders: A systematic review of the clinical research. *Complementary Therapies in Clinical Practice*, 31, 31-37.
9. Armour, M., Smith, C.A., Wang, L.-Q., Naidoo, D., Yang, G.-Y., MacPherson, H., Lee, M.S., & Hay, P. (2019). Acupuncture for depression: A systematic review and meta-analysis. *Journal of Clinical Medicine*, 8(1140), 1-22.
10. Centers for Medicare & Medicaid Services. *National Coverage Determination (NCD) for Acupuncture (30.3)*. CMS website: <https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?NCDId=11&ncdver=1&bc=AAAAGAAAAAAAAA%3d%3d&>.
11. Chen, Z., Wang, Y., Wang, R., Xie, J., & Ren, Y. (2018). Efficacy of acupuncture for treating opioid use disorder in adults: A systematic review and meta-analysis. *Evidence-Based Complementary and Alternative Medicine*, 2018, 3724708, doi.org/10.1155/2018/3724708.

12. Department of Veterans Affairs and Department of Defense (VA/DoD). (2022). Clinical Practice Guidelines for the Management of Major Depressive Disorder. U.S. Department of Veterans Affairs website: <https://www.healthquality.va.gov/guidelines/MH/mdd/VADoDMDDCPGFinal508.pdf>.
13. Deshmukh, S.R., Holmes, J., & Cardno, A. (2018). Art therapy for people with dementia. *Cochrane Database Systematic Review*, 9, CD011073, DOI: 10.1002/14651858.CD011073.pub2.
14. Dunphy, K., Baker, F.A., Dumaresq, E., Carroll-Haskins, K., Eickholt, J., Ercole, M., Kaimal, G., Meyer, K., Sajnani, N., Shamir, O.Y., & Wosch, T. (2019). Creative arts interventions to address depression in older adults: A systematic review of outcomes, processes, and mechanisms. *Frontiers in Psychology*, 9 (2655), 1-24.
15. Fowler, G.C. (2020). Acupuncture. In V. Sierpina (Ed.), Pfenninger & Fowler's procedures for primary care (pp. 4th ed., 1257-1262). Elsevier Publishing.
16. Geipel, J., Koenig, J., Hillecke, T.K., Resch, F., & Kaess, M. (2018). Music-based interventions to reduce internalizing symptoms in children and adolescents: A meta-analysis. *Journal of Affective Disorders*, 225, 647-656.
17. Geretsegger, M., Mössler, K.A., Bieleninik, Ł., Chen, X.J., Heldal, T.O., & Gold, C. (2017). Music therapy for people with schizophrenia and schizophrenia-like disorders. *Cochrane Database Systematic Review*, 5, CD004025, DOI: 10.1002/14651858.CD004025.pub4.
18. Gol, A.S., Ardani, A.R., Farahmand, S.K., Dadgarmoghaddam, M., Ghorani, V., Rezaei, S., & Khorsand, A. (2021). Additive effects of acupuncture in alleviating anxiety: A double-blind, three-arm, randomized clinical trial. *Complementary Therapies in Clinical Practice*, 45(101466), 1-6.
19. Hawkins, E.L., Hawkins, R.D., Dennis, M., Williams, J.M., & Lawrie, S.M. (2019). Animal-assisted therapy for schizophrenia and related disorders: A systematic review. *Journal of Psychiatric Research*, 115, 51-60.
20. Hayes, Inc. (2021). Comparative effectiveness review of acupuncture for the treatment of substance use disorders: A review of reviews. Updated July 29, 2021.
21. Hussain, J.N., Greaves, R.F., & Cohen, M.M. (2019). A hot topic for health: Results of the global sauna survey. *Complementary Therapies in Medicine*, 44, 223-234.
22. Icel, S. and Basogul, C. (2021). Effects of progressive muscle relaxation training with music therapy on sleep and anger of patients at community mental health center. *Complementary Therapies in Clinical Practice*, 43(101338), 1-7.
23. Krause, F., Penzlin, A.I., Ritschel, G., Barlinn, K., Reichmann, H., Weidner, K., Siepman, M., & Siepman, T. (2020). Randomized controlled three-arm study of NADA acupuncture for alcohol addiction. *Addictive Behaviors*, 110(106488), 1-8.
24. Lai, N.M., Chang, S.M.W., Ng, S.S., Tan, S.L., Chaiyakunapruk, N., & Stanaway, F. (2019). Animal-assisted therapy for dementia. *Cochrane Database of Systematic Reviews*, Issue 11, Art. No.: CD013243. DOI: 10.1002/14651858.CD013243.pub2.
25. Lennox, R.D. & Cecchini-Sternquist, M. (2018). Safety and tolerability of sauna detoxification for the protracted withdrawal symptoms of substance abuse. *Journal of International Medical Research*, 46(11), 4480-4499.

26. Li, M., Xing, X., Yao, L., Li, X., He, W., Wang, M., Li, H., Wang, X., Xun, Y., Yan, P., Zhenxing, L., Yang, B., & Yang, K. (2019). Acupuncture for treatment of anxiety, an overview of systematic reviews. *Complementary Therapies in Medicine*, 43(2019), 247-252.
27. Mastrominico, A., Fuchs, T., Manders, E., Steffinger, L., Hirjak, D., Sieber, M., Thomas, E., Holzinger, A., Konrad, A., Bopp, N., & Koch, S.C. (2018). Effects of dance movement therapy on adult patients with autism spectrum disorder: A randomized controlled trial. *Behavioral Sciences*, 8(61), doi:10.3390/bs8070061.
28. Meekums, B. & Karkou, V. (2017). Dance movement therapy for dementia. *Cochrane Database Systematic Review*, 2, CD011022, DOI: 10.1002/14651858.CD011022.pub2.
29. Miller, M., Chen, A.L., Stokes, S.D., Silverman, S., Bowirrat, A., Manka, M., Manka, D., Miller, D.K., Perrine, K., Chen, T.J., Bailey, J.A., Downs, W., Waite, R.L., Madigan, M.A., Braverman, R.R., Damle, U., Mallory, K., Giordano, J., Morse, S., Oscar-Berman, M., Barh, D., & Blum, K. (2012). Early intervention of intravenous KB220IV-neuroadaptagen amino-acid therapy (NAAT) improves behavioral outcomes in a residential addiction treatment program: A pilot study. *Journal of Psychoactive Drugs*, 44(5), 398-409.
30. National Center for Complementary and Integrative Health. (2021). Complementary, alternative, or integrative health: What's in a name? U.S. Department of Health and Human Services. National Institutes of Health website: <https://nccih.nih.gov/health/integrative-health>.
31. Qaseem, A., Barry, M.J., Kansagara, D. for the Clinical Guidelines Committee of the American College of Physicians. (2016). Nonpharmacologic versus pharmacologic treatment of adult patients with major depressive disorder: A clinical practice guideline from the American College of Physicians. *Annals of Internal Medicine*, 164(5), 350-359.
32. Rabeyron, T., Robledo del Canto, J.-P., Carasco, E., Bisson, V., Bodeau, N., Vrait, F.-V., Berna, F., & Bonnot, O. (2020). A randomized controlled trial of 25 sessions comparing music therapy and music listening for children with autism spectrum disorder. *Psychiatry Research*, 293(2020), 113377, 1-7.
33. Salihu, D., Kwan, R., & Wong, E. (2021). The effect of dancing interventions on depression symptoms, anxiety, and stress in adults without musculoskeletal disorders: An integrative review and meta-analysis. *Complementary Therapies in Clinical Practice*, 45(101467), 1-13.
34. Smith, C.A., Armour, M., Lee, M.S., Wang, L.Q., & Hay, P.J. (2018). Acupuncture for depression. *Cochrane Database Systematic Review*, 3, CD004046, DOI: 10.1002/14651858.CD004046.pub4.
35. Srinivasan, S.M., Cavagnino, D.T., & Bhat, A.N. (2018). Effects of equine therapy on individuals with autism spectrum disorder: A systematic review. *Review Journal of Autism and Developmental Disorders*, 5(2), 156-175.
36. Trzmiel, T., Purandare, B., Michalak, M., Zasadzka, E., & Pawlaczyk, M. (2019). Equine assisted activities and therapies in children with autism spectrum disorder: A systematic review and a meta-analysis. *Complementary Therapies in Medicine*, 42, 104-113.
37. van den Noort, M., Yeo, S., Lim, S., Lee, S.H., Staudte, H., & Bosch, P. (2018). Acupuncture as add-on treatment of the positive, negative, and cognitive symptoms of patients with schizophrenia: A systematic review. *Medicines*, 5(29), 1-14.
38. van der Steen, J.T., Smaling, H.J.A., van der Wouden, J.C., Bruinsma, M.S., Scholten, R.J.P.M., & Vink, A.C. (2018). Music-based therapeutic interventions for people with dementia. *Cochrane Database Systematic Review*, CD003477, DOI:10.1002/14651858.CD003477.pub4.

39. White, E., Zippel, J., & Kumar, S. (2020). The effect of equine-assisted therapies on behavioural, psychological and physical symptoms for children with attention deficit/hyperactivity disorder: A systematic review. *Complementary Therapies in Clinical Practice*, 39, 1-10.
40. Zhang, Y., Cai, J., An, L., Hui, F., Ren, T., Hongda, M., & Zhao, Q. (2017). Does music therapy enhance behavioral and cognitive function in elderly dementia patients? A systematic review and meta-analysis. *Ageing Research Reviews*, 35, 1-11.

REVISION HISTORY

Date	Action/Description
12/16/2016	Version 1 (Approved by UMC)
12/16/2017	Annual review performed. Formatting and references updated.
07/15/2019	Annual review performed. Formatting and references updated.
06/15/2020	Annual review: updated references/sourcing.
06/21/2021	Annual review: updated references/sourcing.
06/21/2022	Annual review: updated references/sourcing.

APPENDIX

Additional resources considered in support of this policy:

1. Bidabadi, S.S. & Mehryar, A. (2015). Music therapy as an adjunct to standard treatment for obsessive compulsive disorder and co-morbid anxiety and depression: A randomized clinical trial. *Journal of Affective Disorders*, 184, 13- 17.
2. Geretsegger, M., Elefant, C., Mossler, K.A., & Gold, C. (2014). Music therapy for people with autism spectrum disorder. *Cochrane Database of Systematic Reviews*, 6, CD004381, DOI: 10.1002/14651858.CD004381.pub3.
3. Grant, S., Kandrack, R., Motala, A., Shanman, R., Booth, M., Miles, J., Sorbero, M., & Hempel, S. (2016). Acupuncture for substance use disorders: A systematic review and meta- analysis. *Drug and Alcohol Dependence*, 163, 1-15.
4. Martin, L.A., Koch, S.C., Hirjak, D., & Fuchs, T. (2016). Overcoming disembodiment: The effect of movement therapy on negative symptoms in schizophrenia – a multicenter randomized controlled trial. *Frontiers in Psychology*, 7(483), 1-14.
5. Meekums, B., Karkou, V., & Nelson, E.A. (2015). Dance movement therapy for depression. *Cochrane Database Systematic Review*, 2, CD009895, DOI: 10.1002/14651858.CD009895.pub2.
6. Priebe, S., Savill, M., Wykes, T., Bentall, R.P., Reininghaus, U., Lauber, C., Bremner, S., Eldridge, S., & Rohricht. (2016). Effectiveness of group body psychotherapy for negative symptoms of schizophrenia: Multicenterrandomized controlled trial. *The British Journal of Psychiatry*, 209(1), 54-61.
7. Ren, J. & Xia, J. (2013). Dance therapy for schizophrenia. *Cochrane Database Systematic Review*, 4, DOI: 10.1002/14651858.CD006868.pub3.
8. Uttley, L., Stevenson, M., Scope, A., & Rawdin, A. (2015a). The clinical and cost effectiveness of group art therapy for people with non-psychotic mental health disorders: A systematic review and cost-effectiveness analysis. *BMC Psychiatry*, 19(18), 1-149.
9. Uttley, L., Scope, A., Stevenson, M., Rawdin, A., Taylor Buck, E., Sutton, A., Stevens, J., Kaltenthaler, E., Dent-Brown, K., & Wood, C. (2015b). Systematic review and economic modelling of the clinical effectiveness and cost-effectiveness of art therapy among people with

non-psychotic mental health disorders. *Health Technology Assessment BMC Psychiatry*, 19(18), 1-149.

10. Zhao, K., Bai, Z.G., Bo, A., & Chi, I. (2016). A systematic review and meta-analysis of music therapy for the older adultswith depression. *International Journal of Geriatric Psychiatry*, doi: 10.1002/gps.4494.